

The VIMOS Public Extragalactic Redshift Survey (VIPERS)*

A Support Vector Machine classification of galaxies, stars and AGNs

K. Małek^{†1}, A. Solarz¹, A. Pollo^{2,3}, A. Fritz⁴, B. Garilli^{4,5}, M. Scodeggio⁴, A. Iovino⁶, B. R. Granett⁶, U. Abbas⁷, C. Adami⁵, S. Arnouts^{8,5}, J. Bel⁹, M. Bolzonella¹⁰, D. Bottini⁴, E. Branchini^{11,12,13}, A. Cappi^{10,14}, J. Coupon¹⁵, O. Cucciati¹⁰, I. Davidzon^{10,16}, G. De Lucia¹⁷, S. de la Torre¹⁸, P. Franzetti⁴, M. Fumana⁴, L. Guzzo^{6,19}, O. Ilbert⁵, J. Krywult²⁰, V. Le Brun⁵, O. Le Fevre⁵, D. Maccagni⁴, F. Marulli^{16,21,10}, H. J. McCracken²², L. Paiono⁴, M. Polletta⁴, H. Schlagenhauser^{23,24}, L. A. M. Tasca⁵, R. Tojeiro²⁵, D. Vergani²⁶, A. Zanichelli²⁷, A. Burden²⁵, C. Di Porto¹⁰, A. Marchetti^{28,6}, C. Marinoni⁹, Y. Mellier²², L. Moscardini^{16,21,10}, R. C. Nichol²⁵, J. A. Peacock¹⁸, W. J. Percival²⁵, S. Phleps²⁴, M. Wolk²², and G. Zamorani¹⁰

(Affiliations can be found after the references)

Received -; accepted -;

ABSTRACT

Aims. The aim of this work is to develop a comprehensive method for classifying sources in large sky surveys and we apply the techniques to the VIMOS Public Extragalactic Redshift Survey (VIPERS). Using the optical (u^* , g' , r' , i') and NIR data (z' , K_s), we develop a classifier for identifying stars, AGNs and galaxies improving the purity of the VIPERS sample.

Methods. Support Vector Machine (SVM) supervised learning algorithms allow the automatic classification of objects into two or more classes based on a multidimensional parameter space. In this work, we tailored the SVM for classifying stars, AGNs and galaxies, and applied this classification to the VIPERS data. We train the SVM using spectroscopically confirmed sources from the VIPERS and VVDS surveys.

Results. We tested two SVM classifiers and concluded that including NIR data can significantly improve the efficiency of the classifier. The self-check of the best optical + NIR classifier has shown a 97% accuracy in the classification of galaxies, 97% for stars, and 95% for AGNs in the 5-dimensional colour space. In the test on VIPERS sources with 99% redshift confidence, the classifier gives an accuracy equal to 94% for galaxies, 93% for stars, and 82% for AGNs. The method was applied to sources with low quality spectra to verify their classification, and thus increasing the security of measurements for almost 4 900 objects.

Conclusions. We conclude that the SVM algorithm trained on a carefully selected sample of galaxies, AGNs, and stars outperforms simple colour-colour selection methods, and can be regarded as a very efficient classification method particularly suitable for modern large surveys.

Key words. Cosmology: observations – Methods: statistical – Galaxies: fundamental parameters – Galaxies: active – Galaxies: nuclei – Stars: fundamental parameters

1. Introduction

Over the years, the amount of astronomical data collected by satellites and ground-based surveys is steadily increasing. The zoo of collected data, such as photometry, redshifts, spectral lines, and morphology, is constantly expanding, and increasingly, researchers are turning to automated algorithms to explore

the high-dimensional parameter space. Although computationally challenging, the goal is to make use of every available feature to recognise and extract the most discriminating patterns and allow the full systematisation of the data.

Furthermore, the study of the dependence of galaxy properties on physical parameters such as galaxy mass or environment can greatly benefit from the efficient classification of sources. The classification of different types of sources is one of the basic and at the same time crucial tasks to perform before moving into any scientific analysis.

The first physical classification of sources in a photometric sky survey is between foreground stars within the Galaxy and extragalactic sources. Generally, the distinction between stars and galaxies can be made based upon morphological measurements; point sources are classified as stars while extended sources are classified as galaxies e.g. Vasconcellos et al. (2011); Henrion et al. (2011). At bright apparent magnitudes morphology seems to be a sufficient criterion for classification. However, resolved stellar selection in the current and next generation of wide-field surveys, like Euclid (Laureijs et al. 2012), BigBOSS (Schlegel et al. 2009), DES (Abbott et al. 2005), LSST (Ivezic et al. 2009), LAMOST (Bland-Hawthorn 2012), Pan-STARRS (Kaiser et al. 2010) and/or deep sur-

Send offprint requests to: K. Małek
e-mail: malek.kasia@nagoya-u.jp

*based on observations collected at the European Southern Observatory, Cerro Paranal, Chile, using the Very Large Telescope under programs 182.A-0886 and partly 070.A-9007. Also based on observations obtained with MegaPrime/MegaCam, a joint project of CFHT and CEA/DAPNIA, at the Canada-France-Hawaii Telescope (CFHT), which is operated by the National Research Council (NRC) of Canada, the Institut National des Sciences de l'Univers of the Centre National de la Recherche Scientifique (CNRS) of France, and the University of Hawaii. This work is based in part on data products produced at TERAPIX and the Canadian Astronomy Data Centre as part of the Canada-France-Hawaii Telescope Legacy Survey, a collaborative project of NRC and CNRS. The VIPERS web site is <http://www.vipers.inaf.it/>.

[†]Postdoctoral Fellow of the Japan Society for the Promotion of Science

veys, like VUDS (Le Fèvre et al. 2013, in preparation), HUDF (Beckwith et al. 2006), DLS (Wittman et al. 2002), VISTA (Emerson & Sutherland 2010), is being challenged by the vast number of unresolved galaxies at faint apparent magnitudes (Fadely et al. 2012).

In the case of fainter sources, colour-colour diagrams are the most widely used tools to separate different classes of celestial sources from one another since different types of objects will appear in different colour regions in such diagrams due to the shape of the Spectral Energy Distribution (SED). For example, galaxies possess much redder colours relative to stars due to the higher flux at longer wavelengths (e.g., Walker et al. 1989). Classification methods based on colour-colour selection were employed for star-galaxy separation (e.g. infrared colour diagram used by Pollo et al. 2010) or for finding special classes of sources e.g. high/low redshift quasars, active galactic nuclei, starburst galaxies, or variable stars (Richards et al. 2002; Stern et al. 2005, 2012; Chiu et al. 2005; Brightman & Nandra 2012; Woźniak et al. 2004).

Support Vector Machines (SVMs) are a class of supervised learning algorithms, created as an extension to nonlinear models of the generalised portrait algorithm developed by Vladimir Vapnik (Vapnik 1995), for classification in a multidimensional parameter space. These algorithms are based on the concept of decision planes to classify objects using their relative positions in the n -dimensional parameter space. A large number of observed properties may be analysed simultaneously by the classifier making full use of the data. Within the full parameter space, it is possible to build a more reliable classifier than is possible by only using a subset of the data (for example, by analysing only two photometric colours, instead of the complete set). On the other hand, the method requires a Training Sample, that is, a set of data that have known classifications. Generally, SVM algorithms are sensitive to the measurement errors and are of limited use for extracting information from noisy data sets (Fadely et al. 2012). The classification of observed sources in astronomy is a fundamental problem, and there is still no approach completely free of drawbacks; however SVM algorithms are a novel and very promising classification strategy.

In this paper we apply the SVM algorithm to photometric data. Previous works (e.g., Fadely et al. 2012; Solarz et al. 2012; Vasconcellos et al. 2011; Ball et al. 2006) show high efficiency of that approach for two classes of objects (galaxies and stars). Recently, the Photometric Classification Server (PCS) for the prototype of the Panoramic Survey Telescope and Rapid Response System (Pan-STARRS1) based on Support Vector Machines was developed (Saglia et al., 2012). The PCS system is using five photometric bands: g_{P1} , r_{P1} , g_{P1} , z_{P1} , and y_{P1} , and is able to separate three groups of sources (stars, galaxies, QSOs), without any pre-selection based on colour or redshift range, with high accuracy of galaxy classification ($\sim 97\%$). The purities of stellar and QSO samples' classifications are worse, at the level of 85%, and 83%, respectively.

We decided to develop a three class recognition algorithm, which will be able to classify galaxies/AGNs/stars based on the photometric data in The Canada France Hawaii Telescope Legacy Survey (CFHTLS). We used, as a training set in colour space objects with the best quality spectra from the VIMOS Public Extragalactic Redshift Survey (VIPERS) and VIMOS VLT Deep Survey (VVDS) Deep (F02 field) and Wide (F22 field) data. After careful selection of objects from VIPERS by SVM, and defining characteristic patterns for different types of sources, it will be possible to enlarge the sample of galaxies to be used for more detailed studies. We plan to use this trained

classifier on a large number of sources possessing low quality spectra within VIPERS to recover sources that cannot be classified based upon the spectrum alone. A majority of objects with lower quality spectral information are absorption line systems with low signal-to-noise ratio. Faint red stars and faint passive galaxies are often difficult to distinguish based on their spectral features, if the quality of a spectrum is low. Reconfirmation of a class of such an object by the SVM classifier (galaxy, AGN or star) based upon the photometric measurements increases also the probability that their spectroscopically measured redshift is correct.

The paper is organised as follows. In Section 2, we describe the data used in our analysis, both spectroscopic and photometric. Section 3 describes the principles of the SVM learning algorithm. In Section 4 we introduce the Training Sample used in our work. In Section 5, we compare the efficiency of the classifier with and without near infrared data. Additionally, we present the results of the analysis of the basic tests for the classifiers - self-check and test of the classifier on the VIPERS galaxies with redshift measurements confirmation level equal to 95%. The section closes with the selection of the optimal classifier used for our subsequent analysis. Section 6 describes results of classification of optical near-infrared SVM classifier objects from the VIPERS samples. Finally, in Section 7 we discuss the advantages and limitations of our current SVM classifier and we outline our improvements for the presented classifier.

2. Data

2.1. Photometric data

CFHTLS photometry

The CFHTLS, a joint Canadian-French program, has three distinct survey components: (1) the SuperNovae Legacy Survey the “Deep” survey, (2) the “Wide” - wide synoptic survey (on which VIPERS survey was based), and (3) a very wide shallow survey, the “Very Wide”.

The heart of MegaPrime, the wide-field optical imaging facility, is the MegaCam CCD camera (Boulade et al. 2000). MegaCam, provides multi-colour photometry with wavelength (λ) coverage from 3500 to 9400Å. The main characteristics of the MegaPrime/MegaCam broad band filters are described in Tab. 1. For a more detailed description we refer the reader to the CFHTLS official web page <http://www.cfht.hawaii.edu/Science/CFHTLS/>. All magnitudes used in this paper are the AB photometry system.

Table 1. MegaPrime* and WIRCam** Filters Characteristics.

Filter	u^*	g'	r'	i'	z'	K_s
central λ (nm)	374	487	628	777	1170	2146
bandwidth (nm)	76	145	122	151	687	325
Max. transmission (%)	77.5	93.5	96.3	98	95	98
mag. limit***	25.30	25.50	24.80	24.48	23.60	22.00

* <http://www.cfht.hawaii.edu/Instruments/Filters/megaprime.html>

** <http://www.cfht.hawaii.edu/Instruments/Filters/wircam.html/>

*** measured as the 50% of completeness (MegaPrime) and 5σ (WIRCam) for point sources.

The data used in this work are a part of CFHTLS T0005 release (Mellier et al. 2008), produced at the TERAPIX¹ data centre. We consider a subsample of CFHTLS T0005 catalogue with spectroscopic redshift measured by VIPERS.

WIRCam data

In our work, we also used near-infrared K_s measurements, in the AB magnitude system, corrected for galaxy extinction, taken from Wide-field InfraRed Camera (WIRCam; Thibault et al. 2003; Puget et al. 2004), coming from the dedicated follow-up observations for the VIPERS project (Arnouts et al. 2013, in preparation). The K_s filter has a central wavelength of 2146nm, and maximum transmission on the level of 98%. One may find the detailed description of WIRCam detector on the WIRCam CFHT web page².

2.2. Spectroscopic data

VIPERS survey

The VIMOS Public Extragalactic Redshift Survey (see <http://vipers.inaf.it>) is an ongoing Large Programme aimed at measuring redshifts for $\sim 10^5$ galaxies at redshift $0.5 < z \lesssim 1.2$, to accurately and robustly measure clustering, the growth of structure (through redshift-space distortions) and galaxy properties at an epoch when the Universe was about half its current age. The galaxy target sample is selected from optical photometric catalogues of the Canada-France-Hawaii Telescope Legacy Survey Wide (CFHTLS-Wide, Goranova et al. 2009; Mellier et al. 2008). VIPERS covers $\sim 24 \text{ deg}^2$ on the sky and is divided into two areas within the W1 and W4 CFHTLS fields. Galaxies are selected to a limit of $i_{AB} < 22.5$ measured using SExtractor's `mag_auto` (Kron 1980)-like magnitude. In addition, a simple and robust colour pre-selection in $(g-r)$ vs $(r-i)$ is applied to efficiently remove galaxies at $z < 0.5$. In combination with an efficient observing strategy (Scodeggio et al. 2009), this allows us to double the galaxy sampling rate in the redshift range of interest with respect to a purely magnitude-limited sample, reaching to $\sim 40\%$. At the same time, the area and depth of the survey results in a fairly large volume, $5 \times 10^7 \text{ h}^{-3} \text{ Mpc}^3$, analogous to that of the 2dFGRS at $z \sim 0.1$. This combination of sampling and depth is quite unique over current redshift surveys at $z > 0.5$.

VIPERS spectra are collected with the Visible imaging Multi-Object Spectrograph (VIMOS, Le Fèvre et al. 2000) at moderate resolution ($R = 210$), using the LR Red grism, providing a wavelength coverage of 5500-9500Å, for a typical redshift rms error of $\sigma_z = 0.00047 (1+z)$. The full VIPERS area of $\sim 24 \text{ deg}^2$ is covered through a mosaic of 288 VIMOS pointings (192 in the W1 area, and 96 in the W4 area). Of the VIPERS spectroscopic targets, more than 51 000 K_s counterparts were found: 96% (80%) of our spectra for W1 (W4) field have K_s measurements. More detailed description of WIRCam follow-up survey for VIPERS project can be found in Fritz et al. (2013) and Davidzon et al. (2013).

The redshift quality is quantified at the time of validation by attributing grading flags ($\text{VIPERS}_{Z\text{flag}}$) that are obtained from repeated measurements of redshift for the same sources. The $\text{VIPERS}_{Z\text{flag}}$ for galaxies and stars range from a value of 4, indicating $>99\%$ of confidence that the measurement is secure, to 0,

representing lack of a reliable estimate of redshift. $\text{VIPERS}_{Z\text{flag}}$ equal to 9 correspond to galaxies with only one single clear spectral emission feature. Objects classified as AGNs follow the same scheme but their flags are increased by 10. A similar system was used and tested e.g. for VVDS survey (Le Fèvre et al. 2005). A discussion of the survey data reduction and management infrastructure is presented in Garilli et al. (2012). An early subset of the spectra used here has been analysed and classified through a Principal Component Analysis (PCA) in Marchetti et al. (2012). A more complete description of the survey construction, from the definition of the target sample to the actual spectra and redshift measurements, is given in the parallel survey description paper, Guzzo et al. (2013).

The data set used in this paper are those of the early science data release of VIPERS data as described in Guzzo et al. (2013); see also: de la Torre et al. (2013), Fritz et al. (2013), Marulli et al. (2013), Bel et al. (2013), and Davidzon et al. (2013). This data will be publicly available in fall 2013 as the VIPERS Public Data Release 1 (PDR-1) catalogue. This catalogue includes 55,358 redshifts and corresponds to the reduced data as it was in the VIPERS database at the end of the 2011/2012 observing campaign.

Using the automatic source classifier for VIPERS data is a natural step to handle this unique data volume. Automated and efficient source classifiers based on photometric observations, can provide class labels for catalogues and be used to recover objects for study according to various criteria. Moreover, a multilevel SVM classifier, trained to search for specific types of sources such as Active Galactic Nuclei (AGNs) or galaxies, with an additional redshift measurement as a feature in the parameter space, can be used to boost confidence in the reliability of redshift estimates for sources with poor spectroscopic data. We are planning to develop a more sophisticated and detailed classifier in the near future, enlarging the parameter space by adding measurements of spectral lines and galaxy morphological parameters, thus enabling a finer classification of our sources (e.g. distinguish among different galaxy types).

In this work, we used VIPERS data both to construct a Training Sample and to select samples on which to apply the classifier to separate three different classes of objects (galaxy/AGN/star).

VIMOS-VLT Deep Survey (VVDS)

VIPERS was designed as an extragalactic survey aiming at efficient measuring of redshifts for a large sample of galaxies. To increase the efficiency, stars were carefully removed from the target candidates (which was particularly important for the W4 VIPERS field due to its low galactic latitude). To this aim, both morphological and spectral energy distribution fitting techniques were used (see Guzzo et al. 2013; Coupon et al. 2009). However, it was also important to re-introduce AGNs, which were identified among the stellar objects by their photometric properties (a more detailed description of AGN selection can be found in the survey description paper, Guzzo et al. 2013). Consequently, the number of observed stars and AGNs in VIPERS is quite small.

To construct a reliable Training Sample (see Sec. 3), we included data from another, similar but more complete survey, VVDS. The VVDS fields, like VIPERS, are covered by CFHTLS (and partially by WIRCam observations) and thus the photometric information is homogeneous. Additionally, both surveys utilise the VIMOS spectrograph in similar configurations. The VVDS spectroscopic sample is based upon a purely

¹<http://terapix.iap.fr/>

²<http://www.cfht.hawaii.edu/Instruments/Imaging/WIRCam/>




Fig. 1. An illustration of the operation of the SVM algorithm. The input data (on the left side) are transformed by a kernel into the higher dimensional feature space (right side) where, instead of having a complex boundary separating different classes of objects, we can find an optimal separating hyperplane.

magnitude-limited selection such that the survey contains a much wider variety of sources than VIPERS. We used VVDS-Deep (F02 field) and VVDS-Wide (F22 field) surveys to construct a Training Sample of AGNs (objects classified as AGNs by Gavignaud et al. 2007). The stellar sample was chosen from a part of VVDS Wide F22 which overlaps the VIPERS W4 field.

The Deep F02 survey, covering 0.49 square degrees, is a purely magnitude limited sample to $i_{AB} \leq 24$. The detailed description of the VVDS Deep survey may be found in Le Fèvre et al. (2005). The VVDS Wide F22 survey (Garilli et al. 2008), covering an effective area 3 square degrees, is also a magnitude limited survey with limitation to $i_{AB}=22.5$.

3. Method - Support Vector Machines

The main purpose of the Support Vector Machine (SVM) is to calculate decision planes between a set of objects having different class memberships. A so-called Training Sample, a training set of objects, is used to provide the SVM with examples of the different classes of sources. The SVM searches for the optimal separating hyperplane between the n different classes of objects by maximising the margin between the classes closest points (the so-called support vectors). Instead of using the probability function like in Bayesian statistics or template-fitting methods, the objects are classified based on their relative position in the n -dimensional parameter space with respect to the separation boundary. A well chosen Training Sample is at the heart of the method, because, based on the properties of the Training Sample, the classifier is tuned, and the hyperspace between classes is determined.

The SVM algorithm represents a major development in machine learning techniques. It can be applied to classification or regression problems and is nowadays constantly growing in popularity too to deal with astronomical data to distinguish different classes of sources based on a multidimensional space of parameters taken from observations. Recently, Woźniak et al. (2004) has efficiently used SVMs to analyse variable sources in a 5-dimensional space constructed from the period, amplitude and three colours; Huertas-Company et al. (2008) quantified the morphologies of near-infrared galaxies based on 12-dimensional space, including 5 morphological parameters and other characteristics of galaxies such as luminosity and redshift; Solarz et al. (2012) created a star-galaxy separation algorithm

based on mid and near-infrared colours; Saglia et al. (2012) separated three different classes of sources (galaxies, QSOs, and stars) from PAN-STARRS1 survey, based on five photometric bands. The last year brought a significant number of astronomical papers implementing supervised machine learning algorithms to handle various tasks, not only to classify sources but also to predict characteristic features of specific objects. For example Peng et al. (2012) used SVM to select Active Galactic Nuclei (AGN) candidates and to estimate redshift, Hassan et al. (2013) - to search specific AGN subclass: BL Lacertae and flat-spectrum radio quasars based on the *Second Fermi LAT Catalogue*). Clearly SVMs present an innovative method with a great potential to be widely used in many different branches of astronomy, a potential we are just beginning to tap into.

We use the SVM algorithm to build a non-linear classifier for photometric data to select three different classes of objects: galaxies, AGNs and stars. The first step of our classification task involves selecting a secure Training Sample of galaxies, AGNs and stars, taking advantage of the redshift information provided by VIPERS and VVDS, and using their attributes - i.e. their observed photometric fluxes - to train the SVM.

The algorithm, aided by a non-linear kernel function, searches for a hyperplane which will maximise the distance from the boundary to the closest points belonging to the separate classes of objects (Cristianini & Shawe-Taylor 2000; Shawe-Taylor & Cristianini 2004). The kernel is a symmetric function Φ that maps $k : X \times X \rightarrow F$, so that for all x_i and x_j , $k(x_i, x_j) = \langle \varphi(x_i), \varphi(x_j) \rangle$ from the input space X to the feature space F (see Fig. 1). For our analysis we chose a Gaussian radial basis kernel (RBK) function, defined as:

$$k(x_i, x_j) = \exp(-\gamma \|x_i - x_j\|^2), \quad (1)$$

where $\|x_i - x_j\|$ is the Euclidean distance between x_i , and x_j . The product of the kernel function is a non-linear representation of each parameter from the input to the feature space. The RBK kernel is one of the most popular SVM kernel functions, used to make the non-linear feature map. We decided to use it because of its effectiveness and simplicity in adjusting the free parameters.

A schematic representation of the SVM algorithm classification process is shown in Fig. 2.

For our tasks, we used a soft-boundary SVM method called C-SVM. We chose C-classification because of its good performance and only two free parameters:

- C - a trade-off parameter that sets the width of the margin separating different classes of objects. A large C value sets a small margin of separation between different classes of objects; however increasing the C parameter too much can lead to over-fitting. Reducing C will make the hyperplane between different classes of objects more smooth, allowing for some mis-classifications.
- $\gamma > 0$ parameter (related to the kernel function) determines the topology of the decision surface. A low value of γ sets a very rigid, and complicated decision boundary; a value of γ that is too high can give a very smooth decision surface causing mis-classifications.

For our analysis we used LIBSVM³ (Chang & Lin 2011), an integrated software for support vector classification, which allows for multi-class classification. We used R⁴, a free software environment for statistical computing and graphics, with e1071 interface (Meyer 2001) package installed.

³<http://www.csie.ntu.edu.tw/~cjlin/~libsvm/>

⁴<http://www.r-project.org/>




Fig. 2. A schematic representation of the SVM algorithm classification process. We take as input the pre-selected Training Sample consisting of (in the case of this work) three distinct classes of objects. The SVM is taught how to distinguish one class from the others based on the discriminating properties chosen as feature vectors. Then, the classifier is trained by tuning the free parameters (C and γ). If the result reaches a high enough accuracy rate (the number of objects from the Training Sample that are correctly recognised by the classifier) without overfitting (the resultant hyperplane does not confine too tightly to the sources of a specific type), it will be used to classify the unknown objects (test sample). If the accuracy is not satisfactory, a different parameter space (or Training Sample, if possible) is chosen to tune C and γ . After a number of iterations, which allow the classifier to reach an efficiency level high enough, a real sample can be classified using the discriminant hyperplanes.

4. Training Sample

The successful application of an SVM algorithm requires a carefully selected Training Sample - a set of objects with confirmed classes which will serve as a template for distinguishing the sources whose class we want to determine. Since this work is focused on the selection of galaxies, AGNs, and stars we select as a Training Sample a set of sources whose basic class (galaxy, AGN or star) was established with the highest reliability thanks to their high quality spectra (their redshift being measured with the highest confidence flag within the VIPERS or VVDS surveys). For these sources the accurate photometric information provided by the CFHTLS Wide-survey and the WIRCам follow-up observations of the VIPERS/VVDS fields, provided the colour information needed to create the discriminant vectors for training our SVM algorithm. We produced a model (the optimised C and γ parameters based on the training

data) which predicts the target values of the test data given only the test data attributes (Hsu et al. 2010).

4.1. Galaxies

As a galaxy Training Sample we used the sources with the best redshift measurements in both the W1 and W4 VIPERS fields ($\text{VIPERS}_{\text{Zflag}} = 4$, corresponding to the highest confidence level of redshift measurements and thus of spectroscopic classification as a galaxy). It is useful to remember that VIPERS is pre-selected not only in magnitude ($i' < 22.5$) but also in colours: $(r' - i') > 0.5 * (u^* - g)$ or $(r' - i') > 0.7$. We have divided the galaxy training set into i' -based apparent magnitude-binned samples and trained the classifier on each subset. As a galaxy Training Sample we used 16 271 galaxies: 1 884, 5 483, 6 778, and 3 226 for $19 \leq i' < 20$, $20 \leq i' < 21$, $21 \leq i' < 22$, and $22 \leq i' < 22.5$ apparent magnitude-bins, respectively. Based on our initial tests, we decided to divide our galaxy sample into the magnitude bins to separate more efficiently different groups of galaxies seen in different i' apparent magnitude ranges to improve their classification. Fig. 3 shows that galaxies in different magnitude bins occupy different areas of the colour-colour plots, partially because of different redshift range and different morphology.

4.2. AGNs

Given the small number of AGNs detected in the VIPERS fields with the $\text{VIPERS}_{\text{Zflag}} = 14$, we increased the AGN sample by using all AGNs which had at least 99% confidence level of spectroscopic classification ($\text{VIPERS}_{\text{Zflag}} = 13$ and 14, in total 398 objects). AGN spectra are quite easy to recognize and thus a lower flag on the quality of the measured redshift does not infringe on the reliability of the classification as an AGN. There are two ways that an AGN can be observed in VIPERS:

- it is star-like and meets the AGN candidate selection. This includes samples of X-ray selected AGNs from the XMM-LSS survey, overlapping the VIPERS W1 field (Pierre et al. 2004), and AGNs selected based upon colour-colour criteria from the sample of star-like sources that would otherwise not be targeted.
- it meets the galaxy selection criteria - AGNs which met the galaxy criteria during the main VIPERS colour pre-selection.

We stress that the color pre-selection for galaxies and AGNs is slightly different, and AGNs occupy only a part of the full colour-colour galaxy plane. The first AGN colour separation criterion $\text{CC}_{1\text{AGN}}$:

$$(g' - r') < 1 \wedge \begin{cases} 1. (u^* - g)_{\text{corr}} < 0.6, \\ 2. 0.6 \leq (u^* - g)_{\text{corr}} < 1.2 \& (g' - r')_{\text{corr}} > 0.5(u^* - g')_{\text{corr}} + 0.036, \\ 3. 0.6 \leq (u^* - g)_{\text{corr}} < 2.6 \& (g' - r')_{\text{corr}} < 0.5(u^* - g')_{\text{corr}} + 0.214, \\ 4. (u^* - g')_{\text{corr}} > 2.6, \end{cases} \quad (2)$$

where $(u^* - g)_{\text{corr}}$, and $(g' - r')_{\text{corr}}$ correspond to the color offset. This set of colour-colour selection criteria was inspired from the VVDS survey. After one year of observation it turned out that this allows for stars contamination at the level $\sim 60\%$. From August 2010 additional criterion $\text{CC}_{2\text{AGN}}$, including $(g' - i')$ vs $(u^* - g')$ color-color plane, was added to eliminate stellar sample from AGNs targets. The set of colour-colour criteria included to




Fig. 3. The representative colour-colour plots for galaxy Training Sample. Open black squares represent objects with i' -apparent magnitude between 19 and 20 mag; green X-s - galaxies with i' magnitude between 20 and 21 mag; objects with i' apparent magnitude between $21 \leq i' < 22$, and $22 \leq i' < 22.5$ mag are marked as blue +s and open red triangles, respectively; In the middle panel of colour-colour plots, the boundaries of VIPERS selection are marked as magenta lines.




Fig. 4. The representative colour-colour plots for the AGN Training Sample. Full magenta triangles represent objects brighter than 19 mag in the i' band. Open black triangles - AGNs with i' -apparent magnitude between 19 and 20 mag; open green circles - AGNs with i' magnitude between 20 and 21 mag; objects with i' apparent magnitude between $21 \leq i' < 22$, and $22 \leq i' < 22.5$ mag are marked as open blue squares and open red diamonds, respectively; AGNs with i' apparent magnitude fainter than 22.5 are marked as open rotated cyan triangles.




Fig. 5. The representative colour-colour plots for the star Training Sample. Open black triangles - stars with i' -apparent magnitude between 19 and 20 mag; open green circles - stars with i' magnitude between 20 and 21 mag; objects with i' apparent magnitude between $21 \leq i' < 22$, and $22 \leq i' < 22.5$ mag are marked as open blue squares and open red diamonds, respectively;

CC_{2AGN} is listed below:

$$\begin{cases} 1. (u^* - g')_{corr} < 0.6 \ \& \ -0.2 < (g' - i') < 1, \\ 2. \ 0.6 \leq (u^* - g')_{corr} < 1 \ \& \ -0.2 < (g' - i') < 0.2, \\ 3. (u^* - g')_{corr} \geq 1 \ \& \ (g' - i') < 0.6. \end{cases} \quad (3)$$

Therefore, both criteria (Eq. 2 and Eq. 3) applied simultaneously defined VIPERS AGN targets. However, most of the AGNs share the same colour-colour space as galaxies (as can be seen in Fig. 6), and that is why it is a challenge to distinguish all three classes of objects using an automatic classifier.




Fig. 6. The representative colour-colour plot for VIPERS galaxies with $\text{VIPERS}_{Z_{\text{flag}}} = 4$ (pink x-s) and AGNs with $\text{VIPERS}_{Z_{\text{flag}}} = 3$ and 4 (open blue circles). A part of AGNs occupy different colour-colour area than galaxies and for them, the galaxy/AGN separation is not so difficult. For objects classified as AGNs laying in the same colour-colour plane, the galaxy/AGN/star separation is more challenging. For this reason we decided to use SVM with n -dimensional photometric parameter space to classify sources with similar properties in the typical colour-colour plane.

To enlarge the AGN Training Sample, we also merged the VIPERS sample with objects classified as broad-line AGNs in the VVDS survey. In our Training Sample we included AGNs identified by Gavignaud et al. (2007) - a catalogue of broad emission-line AGNs, from the purely flux-limited spectroscopic sample of the VVDS survey. No color-based pre-selection has been applied in case of these AGNs. For our studies we used 100 AGNs from VVDS Deep F02 (Le Fèvre et al. 2005) and VVDS Wide F22 (Garilli et al. 2008) fields only. We selected these fields since they have the same CFHTLS photometry system as the VIPERS survey.

Cumulatively, our AGN Training Sample reached 498 objects. A part of them, observed by VIPERS, is colour pre-selected. AGNs from VVDS fields have no colour pre-selection (flux-limited only). As we checked on colour-colour plots (see Fig. 4), in the different magnitude bins, we do not see a change in population of our AGN sample with apparent luminosity. For this reason, unlike in the case of the galaxy sample, we decided not to divide the AGN Training Sample into i' -based apparent magnitude binned samples, but to use it as a whole in each bin to increase the population of the training AGNs.

4.3. Stars

VIPERS performed a star/galaxy classification in the CFHTLS wide fields to effectively remove stars from the sample of observed targets. This procedure is crucial since at $i' < 22.5$ the fraction of stars can be as high as 50% (as in the case of W4, Guzzo et al. 2013). The basic VIPERS classification procedure was based on the colour-colour pre-selection with $(r-i) > 0.5 * (u-g)$ or $(r-i) > 0.7$, but due to the low galactic latitude of W4 field, VIPERS implemented an additional procedure. We refer the interested reader to Guzzo et al. (2013) for a complete description on the adopted strategy, here it is suf-

ficient to mention that for objects brighter than $i'=21$ an additional pre-selection based on the observed angular size of sources was applied, while for objects fainter than $i'=21$ a combined method making use of an angular size and SED fitting by the Le Phare code (Arnouts et al. 1999; Ilbert et al. 2006) has been used. These pre-selection criteria proved to be very effective. However, the average stellar contamination in the VIPERS database, for both fields, remains on the level of 3.2% (1.49% and 4.86% for W1 and W4 field, respectively). It means that in the VIPERS PDR-1 catalogue, which includes 55 358 objects, only 1 750 objects have been identified as stars. In sum, the VIPERS PDR-1 catalogue contains 1 750 (3.20%) stars classified as galaxies in the beginning, with colors compatible with an object at $z > 0.5$. This stellar sample can be divided into two main groups:

- stars which were not distinguishable from galaxies based on the VIPERS pre-selection criteria, and
- stars, which were included in the sample as AGN candidates.

Then, it should be stressed that the stars observed by VIPERS are interlopers within the galaxy and AGN samples and thus are not representative of the stellar class. However, our method uses the multi-dimensional colour space which opens a possibility that in such a space, these sources may occupy a region separated from galaxies and AGNs.

To build an unbiased star Training Sample we added spectroscopically classified stars from the VVDS Wide F22 overlap with the VIPERS W4 field. VVDS Wide F22 observations were carried out on the same magnitude limits sample as VIPERS, however, without any photometric pre-selection. The overlap between the VVDS Wide F22 and VIPERS W4 fields contains 920 objects spectroscopically classified as stars by VVDS in the $19 \leq i' < 22.5$ apparent magnitude bin. We increased the stellar Training Sample by using all VIPERS stars with $\text{VIPERS}_{Z_{\text{flag}}} = 4$, in the same apparent magnitude bin (1 312 objects). Cumulatively, our stellar Training Sample reached 2 232 objects.

Similarity to the case of the AGN Training Sample, we did not divide the stellar Training Sample in i' -based apparent magnitude bins. As shown on the representative colour-colour plots for the different magnitude i' bins (Fig. 5), we did not observe a significant change in the distribution of our stellar sample as a function of apparent luminosity.

4.4. Oversampling

Our Training Sample includes more than 16 thousands galaxies, and only 2 232 stars and 498 AGNs (Fig. 7 shows the representative colour-colour plots for galaxies, AGNs and stars chosen for the best Training Sample set). Sampling strategies, such as oversampling and under-sampling, are a popular solution in tackling the problem of classification due to the fact that the SVM classifier is sensitive to a high class imbalance, resulting in a drop in the classification performance (e.g., Tang et al. 2009; Akbani et al. 2004; Raskutti & Kowalczyk 2004). An unbalanced training set tends to over-predict the majority class for unknown sources (Tian et al. 2011).

To avoid this effect, we performed an oversampling of the AGN and stellar training sets so that in each considered magnitude bin we had a similar effective number of objects classified as galaxies, AGNs and stars, respectively. In fact, despite our decision of not splitting AGN and star classes into magnitude bins, unlike we did in the case of galaxies, the imbalance between the numbers of representatives in each class remains high.




Fig. 7. The representative colour-colour plots for all objects used for the Training Sample. Pink x-s represent galaxies. Open blue circles correspond to the AGN sample, and open black squares to the stellar sample.

Table 2. Number (N) of galaxies, AGNs, and stars in our Training Sample after performed oversampling method.

	$19 \leq i' < 20$	$20 \leq i' < 21$	$21 \leq i' < 22$	$22 \leq i' < 22.5$
N Galaxies	1 884	5 483	6 778	2 126
N AGNs	1 520	4 440	5 440	1 760
N Stars	2 232	4 440	5 440	2 232

Using a simple oversampling technique we raised the effective number of AGNs and stars up to $\sim 80\%$ of the number of galaxies in each magnitude bin considered.

We therefore added in each magnitude bin a number of artificial objects calculated as:

$$\lceil X_{i,\text{missing}} \rceil_{10} = N G_i * 0.8 - X \quad (4)$$

where $X_{i,\text{missing}}$ is a number of missing objects (AGNs, stars), and symbol $\lceil \rceil_{10}$ corresponds to rounding the value up to the nearest 10. The additional artificial objects were created by shifting the observed magnitudes by an amount drawn from a Gaussian distribution with $\sigma=0.05$. We also checked how the stellar and AGN Training Samples work if we do not perturb the colours, but instead populate real objects multiple times. As might be expected, the results of classifiers were worse than with randomly modified stars and AGNs.

Tab. 2 summarizes the numbers of training galaxies in each magnitude-binned set together with the number of AGNs and stars after oversampling.

5. Results

5.1. Training procedure

In order to build a classifier which will be able to separate different classes of objects, it is necessary to tune the C and γ parameters using the Training Sample. For the best performance, we have performed a grid-search with values from: $\gamma \in 10^{(-3:-1)}$ and $C \in 10^{(0:3)}$ using a 10-fold cross-validation technique. We first divided the full Training Sample into 10 subsets of equal size and selected 9 subsets to train the classification model and test it against the remaining subset (the so-called self-check). This test was repeated 10 times, with a different subset removed for each training run. The classification accuracy was then averaged over the 10 runs. This process was repeated for each value of the parameters C and γ . In Fig. 8 we present a representative plot of the the grid-search, done for the apparent magnitude




Fig. 8. The mean mis-classification rate as a function of C and γ as estimated from the 10-fold cross-validation technique performed for each pair of parameters (see text for more details). The lower ratio of mis-classification, the better the performance of the SVM algorithm.

bin $19 \leq i' < 20$. The colour of each pointing of the grid codes the mean mis-classification rate of all γ and C values (in log scale on the X and Y axis, respectively). The mis-classification rate is defined as $(1 - \text{Total Accuracy})$ for each magnitude bin (see Eq. 6 further in the paper): the lower the ratio of mis-classification, the better the performance of SVM algorithm. We would like to stress, that a change of the parameter space (such as adding more parameters describing properties of sources) or a sufficient change of the number of training objects inside one class, may result in altering the occupancy of training objects and therefore requires a re-calculation of the best parameters.

To check the efficiency of our classifiers, we count the true objects (true galaxies - TG, true AGNs - TAGN and true stars - TS from the Training Sample originally classified as galaxies, AGNs and stars, respectively), and false objects: FG (false galaxy: when a source from the stellar or AGN Training Sample is classified as a galaxy by the SVM); FS (false star: when an object from a galaxy or AGN Training Sample is classified as a star by the SVM), and FAGN (false AGN: when an object from a galaxy or star Training Sample is classified as an AGN by the SVM). We then calculate the accuracy of our classifier based on the formula:

$$\text{Accuracy} = \frac{\text{TG} + \text{TAGN} + \text{TS}}{\text{TG} + \text{TAGN} + \text{TS} + \text{FG} + \text{FAGN} + \text{FS}}. \quad (5)$$

After completing the 10-fold cross-validation process we calculated the total accuracy of the SVM classifier, defined as the mean accuracy for all iterations:

$$\text{Total Accuracy} = \frac{\sum_{i=0}^N \text{Accuracy}_i}{N}, \quad (6)$$

where $N = 10$ is the number of validation iterations. We performed such a check in each magnitude bin considered.

In our work for galaxy/AGN/star classification we used both a three and five dimensional colour space. The first one was built using only optical data, corresponding to (u^*-g') , $(g'-r)$, and $(r'-i)$ colours, while the second one included NIR data and thus used two extra colours: $(i'-z')$, and $(z'-K_s)$.

5.2. Optical $u^*g'r'i'$ classifier

We constructed colour-colour Training Samples without near-infrared data, based only on the optical u^* , g' , r' , and i' filter bands (a three-dimensional hyperspace). We found that the Total Accuracy as well as the number of correctly classified objects for this approach depend on the apparent magnitude of objects. Averaging over all magnitude bins ($19 \leq i' < 22.5$), once we average results by the number of objects in each bin, the mean Total Accuracy for the optical classifier is equal to 86.39%.

The results of the self-check of our classifier are shown in Tab. 3, showing that only in a few percent of cases (less than 11% in all magnitude bins) galaxies are classified as a star or as an AGN. The most frequent mis-classifications occur in the $19 \leq i' < 20$ bin, in which galaxies are correctly classified at the level of 88.82%, AGNs - 69.45%, and stars at the level of 78.79%. The mis-classifications between stars and galaxies are noticeable in the first three bins. For $20 \leq i' < 21$ and $21 \leq i' < 22$ bins more than 10% of spectroscopically classified stars are classified by the SVM as false galaxies (15.06% and 10.01%, respectively). In the same bins, AGNs are mis-classified as galaxies at the high level of 6.23% and 15.50%, respectively.

The mis-classification of galaxies and AGNs happens mainly in the bins where the percentage of oversampled objects increases. The reason may be related either to our oversampling method or to the worse accuracy of photometry for the fainter sources, as well as to intrinsic properties of classified sources in these bins. We stress that for the SVM method the 100% level of self-check is not desirable since it may be indicative of over-fitting. The boundaries between different classes of objects defined by the Training Sample may become too rigid and artificially complex, not allowing for effective classification of real sources. Nevertheless, it seems that the present, very basic classifier, created on the basis similar to the standard colour-colour approach, works well for our Training Sample.

We next apply our trained classifier to VIPERS galaxies with redshift quality flag $\text{VIPERS}_{\text{Zflag}} = 3$, corresponding to a confidence of the redshift measurements - and correspondingly of correct identification as a galaxy - of $> 99\%$ (hereafter GAL_3). Tab. 5 shows that GAL_3 are correctly classified at a level higher than 85% with an obvious decrease of success with apparent luminosity. The percentage of mis-classifications is constant, at a level of $\sim 10\%$. The strong contamination by false stars is visible for objects fainter than $i'=21$ mag. It is reassuring that this trend is similar to the self-check results (Tab. 3) demonstrating that

Table 5. Test of SVM optical classifier on the galaxies with $\text{VIPERS}_{\text{Zflag}}$ equal to 3. In the first row we show the percentage of correctly classified galaxies. Second and third rows show the percentage of miss-classified galaxies: when a true galaxy is classified by SVM as an AGN or a star, respectively.

	$19 \leq i' < 20$	$20 \leq i' < 21$	$21 \leq i' < 22$	$22 \leq i' < 22.5$
Galaxies	90.97	91.41	85.38	88.82
False AGNs	2.76	2.81	3.06	4.45
False stars	6.27	5.78	11.56	6.73

Table 6. Test of SVM classifier with near-infrared data on the galaxies with $\text{VIPERS}_{\text{Zflag}}$ equal to 3. The first row represents the percentage of correctly classified galaxies. Second and third rows show the percentage of mis-classified galaxies: when a true galaxy is classified by SVM as an AGN or a star, respectively.

	$19 \leq i' < 20$	$20 \leq i' < 21$	$21 \leq i' < 22$	$22 \leq i' < 22.5$
Galaxies	95.38	95.17	93.09	92.72
False AGNs	2.42	2.72	4.30	5.29
False stars	2.20	2.11	2.61	1.99

the Training Sample is representative of the data. In the fainter magnitude bins the photometric errors increase such that the optical u^* , g' , r' , and i' fluxes are not as efficient in distinguishing galaxies and stars.

5.3. Optical+NIR ($u^*g'r'i'z'K_s$) classifier

We enlarge the parameter space by adding the NIR colours (z' and K_s) to our classifier (a five dimensional hyperspace). We performed the same tests as for the optical classifier (self-check, and test on VIPERS GAL_3).

Our Training Sample, composed of exactly the same sources as optical classifier, but with NIR measurements, allows us to train a new optical + NIR classifier. The mean Total Accuracy for this classifier is equal to 94.29%, i.e. higher than the pure optical one. Total Accuracy for particular magnitude bins stays on the similar level $\sim 95\%$ for the whole i' -apparent magnitude binned sample. The constancy of the new classifier for objects fainter than 20 mag in i' band is very promising for the next tests and final classification of VIPERS objects.

Tab. 4 shows the self-check for the u^* , g' , r' , i' , z' and K_s space classifier. Averaging over all magnitude bins, galaxies are correctly classified in $\sim 97.03\%$, AGNs in 95.13%, and stars in 97.05% of the cases, once average results by mean number of objects in each bin. All these numbers are significantly higher than those for purely optical classifier. In case of AGNs, the difference between correctly classified sources for optical and optical+NIR classifiers is equal to 26.46%, 5.46%, 13.30%, and 13.88% for $19 \leq i' < 20$, $20 \leq i' < 21$, $21 \leq i' < 22$, and $22 \leq i' < 22.5$ apparent magnitude-bins, respectively. Stars are correctly classified at higher level than AGNs, with a difference between optical and optical+NIR classifiers equal to 17.58%, 16.79%, 10.94%, and 3.41% for the same magnitude-bins.

Applying this classifier to VIPERS galaxies with $\text{VIPERS}_{\text{Zflag}}$ equal to 3 (GAL_3 , Tab. 6) shows, that galaxies are correctly classified at the mean, very high level of 93.60% (we average results by the number of objects in each bin). Incorrect galaxy classifications, i.e. false AGNs and false stars, are very rare, and do not exceed 2.65% for stars, and 5.30% for AGNs.

Table 3. Results of the self-check of the purely optical classifier (u^* , g' , r' , and i' only). Columns corresponds to the true (spectroscopically classified) galaxies, stars and AGNs. Rows correspond to objects classified as galaxies, AGNs, and stars by our classifier. Then values in bold correspond to the correctly classified objects (galaxies, AGNs, and stars) in defined i' -based apparent magnitude bins. Ratios of classified objects are given in percentage.

	$19 \leq i' < 20$			$20 \leq i' < 21$			$21 \leq i' < 22$			$22 \leq i' < 22.5$		
Total Accuracy	85.01%			87.38%			85.09%			88.09%		
SVM/true	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star
Number of sources	1 884	1 520	2 232	5 483	4 440	4 440	6 778	5 440	5 440	2 126	1 760	2 232
Galaxy	88.82	15.70	10.98	92.10	6.23	15.06	88.39	15.50	10.01	93.18	17.47	3.00
AGN	4.45	69.45	10.23	3.28	90.88	4.48	4.04	81.54	3.81	4.37	79.06	3.28
Star	6.73	14.85	78.79	4.62	2.89	80.46	7.57	2.96	86.19	2.46	3.47	93.72

Table 4. Results of the self-check of the classifier with the near-infrared data (u^* , g' , r' , i' , z' , and K_s). Columns corresponds the true (spectroscopically classified) galaxies, stars and AGNs. Rows correspond to objects classified as galaxies, AGNs, and stars by our classifier. The values marked in bold, are correctly classified objects (galaxies, AGNs, and stars) in defined i' -based apparent magnitude bins. Ratio of classified objects are given in percentage.

	$19 \leq i' < 20$			$20 \leq i' < 21$			$21 \leq i' < 22$			$22 \leq i' < 22.5$		
Total Accuracy	95.47%			95.83%			94.28%			94.58%		
SVM/true	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star
Number of sources	1 884	1 520	2 232	5 483	4 440	4 440	6 778	5 440	5 440	2 126	1 760	2 232
Galaxy	96.28	2.90	1.27	97.61	1.95	0.44	97.11	5.00	2.10	96.10	6.09	1.57
AGN	2.44	95.91	1.70	1.95	96.34	0.80	2.52	94.83	0.77	3.38	92.94	1.30
Star	1.28	1.19	96.37	0.44	0.27	97.25	0.37	0.17	97.13	0.52	0.97	97.13




Fig. 9. Total Accuracy for optical and optical+NIR classifiers (see Tabs. 3 and 4). Results for the optical classifier based on the u^* , g' , r' , and i' filter are marked as a dotted line. Solid line corresponds to the Total Accuracy of the optical+NIR classifier.




Fig. 10. The accuracy of optical and optical+NIR classifiers for VIPERS galaxies with $VIPERS_{zflag}$ equal to 3 (GAL_3). Results for classifier based on the u^* , g' , r' , and i' filter only are marked as a dotted line. Solid line corresponds to the classifier with the NIR data (u^* , g' , r' , i' , z' , and K_s).

We can observe the trend that galaxies have an increased risk of being mis-classified as AGNs in the faintest magnitude bins. One possible explanation of this behaviour is the decrease of the quality of the photometry for the less luminous sources, which have a lower signal-to-noise ratio. On the other hand, the limiting magnitude of CFHTLS is much deeper than the VIPERS one, and photometry should still be fairly good down to $mag\ i' 22.5$. Another explanation could be that some of these galaxies are hosting faint AGNs which were not previously recognized, since with the decreasing luminosity the host galaxy becomes dimmer and the AGN component becomes more significant. This possibility will be further examined in future works.

5.4. Comparison of the classifiers

In Fig. 9 we compare the Total Accuracy for the optical and optical+NIR classifiers. However, on average the classifier based on the u^* , g' , r' , i' , z' and K_s bands, is 7.90% better than the classifier trained without z' and K_s data. Moreover, the Total Accuracy of the optical+NIR classifier decreases very weakly with the apparent magnitude, while such a decrease is slightly stronger for the purely optical classifier; between the first and the second apparent magnitude bin the difference between their Total Accuracy rises from 6.49% to 10.46% from the fainter to the brighter bin.

The preponderance of the classifier constructed with the NIR data is confirmed by the efficiency of correct classification of




Fig. 11. The representative colour-colour plot for all objects used for consistency check for VIPERS object with redshift confirmation level of $> 99\%$, with i' apparent magnitude between 19 and 22.5. Pink x-s represents galaxies with $\text{VIPERS}_{\text{Zflag}}=3$. Open blue circles correspond to AGN sample with redshift confirmation level equal to or higher than 99% ($\text{VIPERS}_{\text{Zflag}}$ equal to 13 and 14). Open black squares correspond to stellar sample with $\text{VIPERS}_{\text{Zflag}}$ equal to 3 and 4.

galaxies with $\text{VIPERS}_{\text{Zflag}}$ equal to 3 (GAL_3). Fig. 10 shows the comparison of Accuracy of both classifiers (with and without near-infrared data) for the GAL_3 sample. For the fainter objects ($21 \leq i' < 22$), the efficiency is decreasing rapidly for the classifier trained without z' and K_s bands, and much smoother for the more sophisticated classifier, trained with infrared features.

We conclude that including NIR data to train the SVM algorithm significantly improves the efficiency of the galaxy/AGN/star classifier. It is evident that NIR features are very important to build an effective classifier for basic astronomical classification these three classes of sources. Based on the above tests, we decided to choose the classifier based on the u^* , g' , r' , i' , z' , and K_s bands to be used in our next analysis.

6. Consistency checks on VIPERS data

6.1. VIPERS objects with redshift confirmation level of $\geq 99\%$

We now apply the optical+NIR classifier to VIPERS data only:

- galaxy sample - all (GAL_3) galaxies in i' -apparent magnitude range between 19 and 22.5 mag, with the total number of sources equal to 13 539,
- AGN sample - all AGNs detected by VIPERS, with redshift confirmation level equal to or higher than 99%, and with i' apparent magnitude between 19 and 22.5 (367 objects). All of these AGNs were used to build the Training Sample (see Sec. 4.2) which means that our classifier should know their position in our 5-dimensional space of parameters, This is not as worrisome as it may look due to the high oversampling needed for AGN sample (more than 200% for the brightest and the faintest apparent magnitude bins, and almost 800% for $20 \leq i' < 21$ and $21 \leq i' < 22$ for i' -apparent magnitude bins) that significantly erases the possibly peculiar characteristics of the 367 AGN chosen for the Training Sample.

- stellar sample - all spectroscopically detected stars, with confirmation level of $> 99\%$ ($\text{VIPERS}_{\text{Zflag}}$ equal to 3 and 4), and i' apparent magnitude between 19 and 22.5 (1 729 stars). All stars with $\text{VIPERS}_{\text{Zflag}}=4$ were used as a part of stellar Training Sample.

Fig. 11 shows the representative colour-colour plot for GAL_3 , AGNs with $\text{VIPERS}_{\text{Zflag}}$ equal to 13 and 14, and stars with $\text{VIPERS}_{\text{Zflag}}$ equal to 3 and 4, chosen for the consistency check.

For this test, all three classes of sources were divided into four i' -apparent magnitude bins ($19 \leq i' < 20$, $20 \leq i' < 21$, $21 \leq i' < 22$, and $22 \leq i' < 22.5$), the same as used in the Training Sample. Then, we applied our optical+NIR classifier on this data.

Tab. 7 shows the results of the automatic classification.

The mean accuracy for galaxies, averaged over the mean number of objects in each apparent magnitude bin, equals 93.60%. This result for galaxy classification displays only slightly less level of efficiency ($\sim 1.50\%$) than galaxy classification obtained during the self-check of the classifier (see Sec. 5.3). It means that the hyperspace of galaxy parameters used for the Training Sample is well defined.

The result of AGN classification is worse than the one obtained during the self-check but still satisfactory. Averaging over all magnitude bins, AGNs are correctly classified at a level equal to 81.80% with a significant decrease with i' apparent magnitude between 21 and 22 mag. Stars are correctly classified at the high mean level of 92.52% with a significant drop for the $22 \leq i' < 22.5$ apparent magnitude bin (84.47%). The performance of the classifier in the case of AGNs may look relatively poor. However, as already mentioned, we should keep in mind that the VIPERS selection allows AGNs pre-classified as galaxies or stars based on their colour properties. Keeping this in mind, we should rather feel satisfied that a high fraction of these AGNs can be separated into a different section of the 5-dimensional hyperspace from galaxies and stars, using AGN Training Sample consists only 498 objects.

We did not find any crucial mis-classifications for the galaxy sample. The galaxies are classified correctly, on a very high level. For the AGN sample, the contamination of true AGNs classified as a galaxies (8.17%, 7.37%, 10.46%, 14.90% for the $19 \leq i' < 20$, $20 \leq i' < 21$, $21 \leq i' < 22$, and $22 \leq i' < 22.5$ bins, respectively) and stars (8.96%, 10.55%, 6.79%, 9.56% for the $19 \leq i' < 20$, $20 \leq i' < 21$, $21 \leq i' < 22$, and $22 \leq i' < 22.5$ bins, respectively) is significant. For the stellar sample, the classifier more often mis-classified true stars as galaxies than AGNs. In the future development of this classifier, we will include the morphological information as well as emission/absorption lines, which should improve the algorithm and increase the percentage of correctly classified sources as well.

6.2. VIPERS objects with redshift confirmation level lower than 99%

We preformed a classification for VIPERS objects with confirmation level lower than 99%. In particular, we used galaxies, AGNs, and stars from the VIPERS database, with quality of the measured redshift, $\text{VIPERS}_{\text{Zflag}}$, equal to 2 and 1. $\text{VIPERS}_{\text{Zflag}}$ equals to 2 means that the measured redshift is fairly secure, with confidence level $\geq 95\%$. Objects with $\text{VIPERS}_{\text{Zflag}}$ equal to 1 are more tentative, and their redshift measurement was based on weak spectral features and/or continuum shape. For these objects there is a $\sim 50\%$ probability that the redshift could be wrong. A more detailed description of the $\text{VIPERS}_{\text{Zflag}}$ and qual-

Table 7. Results of the test of the optical+NIR classifier for GAL₃, and AGNs and stars with redshifts measurements on a confirmation level $\geq$ to 99%. AGNs from this sample were used for building Training Sample. The rest of the objects is not related to Training Sample. Values marked in bold correspond to the correctly classified objects (galaxies, AGNs, and stars) in i' -based apparent magnitude bins. The ratio of the classified objects is given in percentage.

	19 $\leq i' < 20$			20 $\leq i' < 21$			21 $\leq i' < 22$			22 $\leq i' < 22.5$		
SVM/true	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star
Number of sources	445	69	337	3 271	1340	428	7 667	127	701	2 156	37	263
Galaxy	95.38	12.52	4.17	95.17	7.37	3.27	93.09	10.46	3.42	92.72	14.90	9.09
AGN	2.42	77.34	3.70	2.72	82.08	3.27	4.30	82.75	1.43	5.29	75.54	6.44
Star	2.20	10.14	92.13	2.11	10.55	93.46	2.61	6.79	95.15	1.99	9.56	84.47

Table 8. Results of the optical + NIR classifier for galaxies, AGNs, and stars with redshifts measurements on a confirmation level equal to 95% (VIPERS_{Zflag} = 2). Objects are not related to Training Sample. Values marked in bold correspond to the correctly classified objects (galaxies, AGNs, and stars) in i' -based apparent magnitude bins. The ratio of the classified objects is given in percentage.

	19 $\leq i' < 20$			20 $\leq i' < 21$			21 $\leq i' < 22$			22 $\leq i' < 22.5$		
SVM/true	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star
Number of sources	8	33	10	945	75	27	5 757	80	145	6 226	48	159
Galaxy	84.15	6.25	30.00	94.18	12.00	22.22	92.81	20.25	29.65	58.62	4.17	17.61
AGN	9.75	93.75	40.00	3.49	88.00	29.63	4.41	77.22	11.03	20.56	93.75	18.87
Star	6.10	0.00	30.00	2.33	0.00	48.15	2.78	2.53	59.32	20.82	2.08	63.52

Table 9. The results of the optical + NIR classifier for galaxies, AGNs, and stars with redshifts measurements on a confirmation level equal to 50% (VIPERS_{Zflag} equals to 1). Objects are not connected with Training Sample. Values marked in bold correspond to the correctly classified objects (galaxies, AGNs, and stars) in i' -based apparent magnitude bins. The ratio of the classified objects is given in percentage.

	19 $\leq i' < 20$			20 $\leq i' < 21$			21 $\leq i' < 22$			22 $\leq i' < 22.5$		
SVM/true	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star	Galaxy	AGN	Star
Number of sources	35	8	4	355	13	24	2 833	35	81	3 157	30	139
Galaxy	85.71	50.00	25.00	92.68	23.08	37.50	88.14	20.00	60.56	43.09	23.33	25.90
AGN	11.43	50.00	75.00	3.94	76.92	37.50	7.12	68.57	9.86	31.03	66.67	31.65
Star	2.86	0.00	0.00	3.38	0.00	25.00	4.74	11.43	29.58	25.88	10.00	42.45

ity of measured redshifts can be found in Guzzo et al. (2013), and Garilli et al. (2012).

Results of SVM classification of objects with VIPERS_{Zflag} equal to 2 (Tab. 8) and 1 (Tab. 9) show a very good conformity to the previously user supervised estimations. Galaxies are classified with agreement to redshift measurements on the mean level of 76.45% for VIPERS_{Zflag} = 2 and 66.08% for VIPERS_{Zflag} = 1.

The ongoing scientific analysis of galaxy evolution and clustering is mainly based on objects which have a secure redshift measurements (VIPERS_{Zflag} $\geq$ 2, depending on the topic). With the SVM classification we can reconfirm the identify of galaxies with the lower quality flags and thus increase the number of galaxies which could be used for more detailed analysis. This may apply to 4 735 galaxies, 58 AGNs, and 86 stars with VIPERS_{Zflag} equal to 1 (these numbers were calculated as a sum of galaxies, AGNs, and stars which were classified to the same class of objects during redshift validation and by optical+NIR classifier; marked in bold in Tab. 9). This method may also reconfirm the class to 9 952 galaxies, 177 AGNs and 160 stars with VIPERS_{Zflag} = 2 classified as galaxies, AGNs, and stars by checking the results twice by different observers, and by our classifier.

One may argue that the VIPERS_{Zflag} is related to the redshift value, not the identification of the galaxy itself. However, it

should be noted that a majority of sources with low VIPERS_{Zflag} are absorption line systems with noisy, low signal-to-noise spectra. Galaxies with such spectra can be particularly easily misclassified as stars during the spectroscopic measurement process, either automatic, or human-supervised. For instance, typical features of the elliptical galaxy at $z \sim 1$, around Balmer break, can be confused with characteristic features of the M-type star. In such cases, an independent confirmation that the position of an object in the 5-dimensional colour space is actually typical for a galaxy, actually increases also probability that its redshift has been assigned correctly.

The number of stars and AGNs in the sample of objects with VIPERS_{Zflag} equal to 1 and 2 is very low (a few objects in the brightest apparent luminosity bin, and a few dozens for objects with i' magnitude lower than 21 mag). This fact results from an initial star/galaxy separation performed by VIPERS. In the VIPERS database the stars, which remained after the colour-colour pre-selection, are not typical, and they occupy a similar area as galaxies on the colour-colour plots. Then, the fact that we can reconfirm the identity of a significant fraction of them, can be already regarded as a success. The spectra of stars with VIPERS_{Zflag} = 1 or 2, which were classified as galaxies by our classifier, will have to be re-examined since some of them might be also be genuine galaxies.

We should consider the fact that VIPERS galaxy sample includes different types of AGNs, which are in the VIPERS database classified as galaxies, also for objects with $\text{VIPERS}_{\text{Zflag}}$ higher than 3. During the standard redshift measurements process only the broad line AGNs are being recognized and flagged. It implies that our galaxy Training Sample contains, in addition to a pure sample of normal galaxies, also specific types of AGNs, otherwise difficult to recognize in VIPERS spectra during a standard redshift measurements process. VIPERS *hunt* for the specific types of AGNs lurking within the heap of collected sources is still ongoing, hence, at least for now, we are forced to work with the data composed of both galaxies and AGNs. The contamination of galaxies and AGNs is most prominent for the faintest bin ($22 \leq i' < 22.5$), where more than 20% (30%) of objects classified as galaxies with $\text{VIPERS}_{\text{Zflag}}$ equal to 2 (1) are identified as AGNs by optical+NIR SVM classifier.

We look forward to use SVM methods adding further information on spectral lines and source morphologies as a very promising tool to improve classification for fainter sources and to refine further classes of objects that the software can discriminate.

7. Conclusions

Application of the Support Vector Machine algorithm can deliver an excellent (with accuracy level for self-check test higher than 98% for galaxies, 94% for AGNs, and 93% for stars) classification for three classes of objects, after a careful selection of the Training Sample. For our analysis we constructed two classifiers, with and without near infrared data using a multi-dimensional colour hyperspace. A part of the AGN and star samples were extracted from the VVDS survey. We have found a significant improvement of the SVM classification (8% in the Total Accuracy of classifier) adding a NIR colour parameters to our feature vectors.

For the optical+NIR classifier we obtained a very good agreement (93.60%, 81.80%, and 92.52% for galaxies, AGNs, and stars, respectively) with the VIPERS spectroscopic sample with flag confidence level of z measurements equal to 95%. What makes our approach to SVM classification exceptional is that due to the enormous amount of excellent quality data we could create the classifier which was trained on the part of the most secure sources and then test it against the remaining secure objects to create the most efficient pattern recognition system. The VIPERS survey gathered a large number of sources (55 358) with very good spectroscopic measurements, which then were strictly analysed to obtain the most secure redshifts. This fact allowed for the choice of the best sample, which could be used as a base for the new method of automatic classification.

SVM classifiers are mostly used in the literature for separation of two classes of sources (e.g. stars and galaxies). The only recent application of the SVM to the galaxy/AGNs/stars classification was performed by Saglia et al. (2012), who trained and used his classifier for the Pan-STARRS1 data. Comparing the accuracies of our classifier and those of Saglia et al. (2012) we found that our self-check results look somewhat better (97%, 95%, 97% vs 97%, 84%, 85% for galaxies, AGNs, stars for VIPERS and Pan-STARRS1 classifier, respectively). However, we have to stress that both methods cannot be directly compared because of initial differences in both surveys. Pan-STARRS1 is a purely limited survey, which implies a much higher variety of properties of all the sources it contains. In contrary, VIPERS was pre-selected to contain only $0.5 < z < 1.2$ galaxies, which assures that they form a much more distinct and better separated group

in a multi-colour space. This may facilitate a separation between galaxies and AGNs, as well as a part of stars which were re-introduced to the VIPERS target sample as AGN candidates. On the other hand, the lack of ‘typical’ stars in the VIPERS database (rejected after colour, and half-light radius pre-selection) occupying the same colour-colour space as galaxies, may hamper our classification based only on colours, and decrease the efficiency of our classifier for sources from the real sample.

Our approach allows us to photometrically classify sources in the VIPERS survey, augmenting the spectral information. By classifying the sources with low-quality spectra, we can improve the classification, enlarging the samples that may be used for analysis. Using the optical+NIR classifier, we have confirmed the class of 4 900 objects with low flag. Further improvement of our classifier by the addition of the morphology and emission/absorption line information will improve the already very good performance of galaxy/AGN/star classifier. It will also allow for the development of a more specific galaxy and AGN types classifications.

Acknowledgements. We acknowledge the crucial contribution of the ESO staff for the management of service observations. In particular, we are deeply grateful to M. Hilker for his constant help and support of this program. Italian participation to VIPERS has been funded by INAF through PRIN 2008 and 2010 programs. LG and BRG acknowledge support of the European Research Council through the Darklight ERC Advanced Research Grant (# 291521). OLF acknowledges support of the European Research Council through the EARLY ERC Advanced Research Grant (# 268107). Polish participants have been supported by the Polish Ministry of Science (grant N N203 51 29 38), the Polish-Swiss Astro Project (co-financed by a grant from Switzerland, through the Swiss Contribution to the enlarged European Union), the European Associated Laboratory Astrophysics Poland-France HECOLS and a Japan Society for the Promotion of Science (JSPS) Postdoctoral Fellowship for Foreign Researchers (KM, P11802). AS has been supported by the Global COE Program Request for Fundamental Principles in the Universe: from Particles to the Solar System and the Cosmos commissioned by the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan, and by the JSPS Strategic Young Researcher Overseas Visits Program for Accelerating Brain Circulation, Construction of a Global Platform for the Study of Sustainable Humanosphere. GDL acknowledges financial support from the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement n. 202781. WJP and RT acknowledge financial support from the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007-2013)/ERC grant agreement n. 202686. WJP is also grateful for support from the UK Science and Technology Facilities Council through the grant ST/I001204/1. EB, FM and LM acknowledge the support from grants ASI-INAf I/023/12/0 and PRIN MIUR 2010-2011. YM acknowledges support from CNRS/INSU (Institut National des Sciences de l’Univers) and the Programme National Galaxies et Cosmologie (PNCG). CM is grateful for support from specific project funding of the *Institut Universitaire de France* and the LABEX OCEVU.

References

- Abbott, T., Aldering, G., Annis, J., et al. 2005, ArXiv Astrophysics e-prints
- Akbani, R., Kwek, S., & Japkowicz, N. 2004, in In Proceedings of the 15th European Conference on Machine Learning (ECML), 39–50
- Arnouts, S., Cristiani, S., Moscardini, L., et al. 1999, MNRAS, 310, 540
- Ball, N. M., Brunner, R. J., Myers, A. D., & Tchong, D. 2006, ApJ, 650, 497
- Beckwith, S. V. W., Stiavelli, M., Koekemoer, A. M., et al. 2006, AJ, 132, 1729
- Bel, J. et al. 2013, A&A, , submitted
- Bland-Hawthorn, J. 2012, Research in Astronomy and Astrophysics, 12
- Boulade, O., Charlot, X., Abbon, P., et al. 2000, in Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, Vol. 4008, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. M. Iye & A. F. Moorwood, 657–668
- Brightman, M. & Nandra, K. 2012, MNRAS, 422, 1166
- Chang, C.-C. & Lin, C.-J. 2011, ACM Transactions on Intelligent Systems and Technology, 2, 27:1, software available at <http://www.csie.ntu.edu.tw/~cjlin/libsvm>, issue = 3
- Chiu, K., Zheng, W., Schneider, D. P., et al. 2005, AJ, 130, 13
- Coupon, J., Ilbert, O., Kilbinger, M., et al. 2009, A&A, 500, 981

- Cristianini, N. & Shawe-Taylor, J. 2000, *An Introduction to Support Vector Machines: And Other Kernel-Based Learning Methods* (Cambridge University Press)
- Davidzon, I., Bolzonella, M., et al. 2013, *A&A*, submitted, ArXiv e-print 1303.3808
- de la Torre, S., Guzzo, L., Peacock, J. A., et al. 2013, *A&A*, submitted, ArXiv e-print 1303.2622
- Emerson, J. & Sutherland, W. 2010, *The Messenger*, 139, 2
- Fadely, R., Hogg, D. W., & Willman, B. 2012, *ApJ*, 760, 15
- Fritz, A., Scoddeggio, M., et al. 2013, *A&A*, submitted
- Garilli, B., Le Fèvre, O., Guzzo, L., et al. 2008, *A&A*, 486, 683
- Garilli, B., Paiono, L., Scoddeggio, M., et al. 2012, *PASP*, 124, 1232
- Gavignaud, I., Bongiorno, A., Paltani, S., et al. 2007, *VizieR Online Data Catalog*, 345, 70079
- Goranova, Y., Hudelot, P., Contini, T., et al. 2009, *The CFHTLS T0006 Release*, http://terapix.iap.fr/cpl/table_syn_T0006.html
- Guzzo, L., Scoddeggio, M., Garilli, B., et al. 2013, *A&A*, submitted, ArXiv e-print 1303.2623
- Hassan, T., Mirabal, N., Contreras, J. L., & Oya, I. 2013, *MNRAS*, 428, 220
- Henrion, M., Mortlock, D. J., Hand, D. J., & Gandy, A. 2011, *MNRAS*, 412, 2286
- Hsu, C.-W., Chang, C. C., & C.-J., L. 2010, *A Practical Guide to Support Vector Classification*, Department of Computer Science, National Taiwan University, Taiwan, <http://www.csie.ntu.edu.tw/~cjlin/papers/guide/guide.pdf>
- Huertas-Company, M., Rouan, D., Tasca, L., Soucaill, G., & Le Fèvre, O. 2008, *A&A*, 478, 971
- Ilbert, O., Arnouts, S., McCracken, H. J., et al. 2006, *A&A*, 457, 841
- Ivezic, Z., Tyson, J. A., Axelrod, T., et al. 2009, in *Bulletin of the American Astronomical Society*, Vol. 41, American Astronomical Society Meeting Abstracts 213, 460.03
- Kaiser, N., Burgett, W., Chambers, K., et al. 2010, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 7733, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series
- Kron, R. G. 1980, *ApJS*, 43, 305
- Laureijs, R., Gondoin, P., Duvet, L., et al. 2012, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 8442, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series
- Le Fèvre, O., Saisse, M., Mancini, D., et al. 2000, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 4008, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. M. Iye & A. F. Moorwood, 546–557
- Le Fèvre, O., Vettolani, G., Garilli, B., et al. 2005, *A&A*, 439, 845
- Le Fèvre, O. et al. 2013, in preparation, <http://cesam.oamp.fr/vuds>
- Marchetti, A., Granett, B. R., Guzzo, L., et al. 2012, *MNRAS*, 107
- Marulli, F., Bolzonella, M., Branchini, E., et al. 2013, *A&A*, submitted, ArXiv e-print 1303.2633
- Mellier, Y., Bertin, E., Hudelot, P., et al. 2008, *The CFHTLS T0005 Release*, <http://terapix.iap.fr/cpl/oldSite/Descart/CFHTLS-T0005-Release.pdf>
- Meyer, D. 2001, *R News*, 1, 23
- Peng, N., Zhang, Y., Zhao, Y., & Wu, X.-b. 2012, *MNRAS*, 425, 2599
- Pierre, M., Valtchanov, I., Altieri, B., et al. 2004, *J. Cosmology Astropart. Phys.*, 9, 11
- Pollo, A., Rybka, P., & Takeuchi, T. T. 2010, *A&A*, 514, A3
- Puget, P., Stadler, E., Doyon, R., et al. 2004, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 5492, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. A. F. M. Moorwood & M. Iye, 978–987
- Raskutti, B. & Kowalczyk, A. 2004, *SIGKDD Explor. Newsl.*, 6, 60
- Richards, G. T., Fan, X., Newberg, H. J., et al. 2002, *AJ*, 123, 2945
- Saglia, R. P., Tonry, J. L., Bender, R., et al. 2012, *ApJ*, 746, 128
- Schlegel, D. J., Bebek, C., Heetderks, H., et al. 2009, *ArXiv e-prints*
- Scoddeggio, M., Franzetti, P., Garilli, B., Le Fèvre, O., & Guzzo, L. 2009, *The Messenger*, 135, 13
- Shawe-Taylor, J. & Cristianini, N. 2004, *Kernel Methods for Pattern Analysis* (Cambridge University Press)
- Solarz, A., Pollo, A., Takeuchi, T. T., et al. 2012, *A&A*, 541, A50
- Stern, D., Assef, R. J., Benford, D. J., et al. 2012, *ApJ*, 753, 30
- Stern, D., Eisenhardt, P., Gorjian, V., et al. 2005, *ApJ*, 631, 163
- Tang, Y., Zhang, Y.-Q., Chawla, N. V., & Krasser, S. 2009, *IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics)*, 39, 281
- Thibault, S., Cui, Q., Poirier, M., et al. 2003, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 4841, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. M. Iye & A. F. M. Moorwood, 932–943
- Tian, J., Gu, H., & Liu, W. 2011, *Neural Comput. Appl.*, 20, 203
- Vapnik, V. N. 1995, *The Nature of Statistical Learning Theory* (Springer)
- Vasconcellos, E. C., de Carvalho, R. R., Gal, R. R., et al. 2011, *AJ*, 141, 189
- Walker, H. J., Volk, K., Wainscoat, R. J., Schwartz, D. E., & Cohen, M. 1989, *AJ*, 98, 2163
- Wittman, D. M., Tyson, J. A., Dell’Antonio, I. P., et al. 2002, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 4836, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, ed. J. A. Tyson & S. Wolff, 73–82
- Woźniak, P. R., Williams, S. J., Vestrand, W. T., & Gupta, V. 2004, *AJ*, 128, 2965
- 1 Department of Particle and Astrophysical Science, Nagoya University, Furo-cho, Chikusa-ku, 464-8602 Nagoya, Japan
 - 2 Astronomical Observatory of the Jagiellonian University, Orla 171, 30-001 Cracow, Poland
 - 3 National Centre for Nuclear Research, ul. Hoza 69, 00-681 Warszawa, Poland
 - 4 INAF - Istituto di Astrofisica Spaziale e Fisica Cosmica Milano, via Bassini 15, 20133 Milano, Italy
 - 5 Aix Marseille Université, CNRS, LAM (Laboratoire d’Astrophysique de Marseille) UMR 7326, 13388, Marseille, France
 - 6 INAF - Osservatorio Astronomico di Brera, Via Brera 28, 20122 Milano, via E. Bianchi 46, 23807 Merate, Italy
 - 7 INAF - Osservatorio Astrofisico di Torino, 10025 Pino Torinese, Italy
 - 8 Canada-France-Hawaii Telescope, 65–1238 Mamalahoa Highway, Kamuela, HI 96743, USA
 - 9 Aix-Marseille Université, CNRS, CPT (Centre de Physique Théorique) UMR 7332, F-13288 Marseille, France
 - 10 INAF - Osservatorio Astronomico di Bologna, via Ranzani 1, I-40127, Bologna, Italy
 - 11 Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, via della Vasca Navale 84, 00146 Roma, Italy
 - 12 INFN, Sezione di Roma Tre, via della Vasca Navale 84, I-00146 Roma, Italy
 - 13 INAF - Osservatorio Astronomico di Roma, via Frascati 33, I-00040 Monte Porzio Catone (RM), Italy
 - 14 Laboratoire Lagrange, UMR7293, Université de Nice Sophia-Antipolis, CNRS, Observatoire de la Côte d’Azur, 06300 Nice, France
 - 15 Institute of Astronomy and Astrophysics, Academia Sinica, P.O. Box 23-141, Taipei 10617, Taiwan
 - 16 Dipartimento di Fisica e Astronomia - Università di Bologna, viale Berti Pichat 6/2, I-40127 Bologna, Italy
 - 17 INAF - Osservatorio Astronomico di Trieste, via G. B. Tiepolo 11, 34143 Trieste, Italy
 - 18 SUPA, Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, United Kingdom
 - 19 Dipartimento di Fisica, Università di Milano-Bicocca, P.zza della Scienza 3, I-20126 Milano, Italy
 - 20 Institute of Physics, Jan Kochanowski University, ul. Światokrzyska 15, 25-406 Kielce, Poland
 - 21 INFN, Sezione di Bologna, viale Berti Pichat 6/2, I-40127 Bologna, Italy
 - 22 Institute d’Astrophysique de Paris, UMR7095 CNRS, Université Pierre et Marie Curie, 98 bis Boulevard Arago, 75014 Paris, France
 - 23 Universitätssternwarte München, Ludwig-Maximilians Universität, Scheinerstr. 1, D-81679 München, Germany
 - 24 Max-Planck-Institut für Extraterrestrische Physik, D-84571 Garching b. München, Germany
 - 25 Institute of Cosmology and Gravitation, Dennis Sciama Building, University of Portsmouth, Burnaby Road, Portsmouth, PO1 3FX, United Kingdom
 - 26 INAF - Istituto di Astrofisica Spaziale e Fisica Cosmica Bologna, via Gobetti 101, I-40129 Bologna, Italy
 - 27 INAF - Istituto di Radioastronomia, via Gobetti 101, I-40129, Bologna, Italy
 - 28 Università degli Studi di Milano, via G. Celoria 16, 20130 Milano, Italy
 - 29 Institut Universitaire de France, 103, bd. Saint-Michel, F-75005 Paris, France